



**Sensing and regulation of cell volume - we know so much and yet understand so little  
TRPV4 as a sensor of volume changes but possibly without a volume-regulatory role?**

Toft-Bertelsen, Trine L.; Larsen, Brian R.; MacAulay, Nanna

*Published in:*  
Channels (Austin)

*DOI:*  
[10.1080/19336950.2018.1438009](https://doi.org/10.1080/19336950.2018.1438009)

*Publication date:*  
2018

*Document version*  
Publisher's PDF, also known as Version of record

*Document license:*  
[CC BY](#)

*Citation for published version (APA):*  
Toft-Bertelsen, T. L., Larsen, B. R., & MacAulay, N. (2018). Sensing and regulation of cell volume - we know so much and yet understand so little: TRPV4 as a sensor of volume changes but possibly without a volume-regulatory role? *Channels (Austin)*, 12(1), 100-108. <https://doi.org/10.1080/19336950.2018.1438009>



## Sensing and regulation of cell volume – we know so much and yet understand so little: TRPV4 as a sensor of volume changes but possibly without a volume-regulatory role?

Trine L. Toft-Bertelsen, Brian R. Larsen & Nanna MacAulay

To cite this article: Trine L. Toft-Bertelsen, Brian R. Larsen & Nanna MacAulay (2018) Sensing and regulation of cell volume – we know so much and yet understand so little: TRPV4 as a sensor of volume changes but possibly without a volume-regulatory role?, Channels, 12:1, 100-108, DOI: [10.1080/19336950.2018.1438009](https://doi.org/10.1080/19336950.2018.1438009)

To link to this article: <https://doi.org/10.1080/19336950.2018.1438009>



© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Accepted author version posted online: 09 Feb 2018.  
Published online: 09 Feb 2018.



Submit your article to this journal [↗](#)



Article views: 272



View related articles [↗](#)



View Crossmark data [↗](#)

## Sensing and regulation of cell volume – we know so much and yet understand so little: TRPV4 as a sensor of volume changes but possibly without a volume-regulatory role?

Trine L. Toft-Bertelsen , Brian R. Larsen , and Nanna MacAulay 

Department of Neuroscience, University of Copenhagen, Copenhagen, Denmark

### ABSTRACT

Cellular volume changes lead to initiation of cell volume regulatory events, the molecular identity of which remains unresolved. We here discuss experimental challenges associated with investigation of volume regulation during application of large, non-physiological osmotic gradients. The TRPV4 ion channel responds to volume increase irrespectively of the molecular mechanism underlying cell swelling, and is thus considered a sensor of volume changes. Evidence pointing towards the involvement of TRPV4 in subsequent volume regulatory mechanisms is intriguing, yet far from conclusive. We here present an experimental setting with astrocytic cell swelling in the absence of externally applied osmotic gradients, and the lack of evidence for involvement of TRPV4 in this regulatory volume response. Our aim with these new data and the preceding discussion is to stimulate further experimental effort in this area of research to clarify the role of TRPV4 and other channels and transporters in regulatory volume responses.

### ARTICLE HISTORY

Received 21 December 2017  
Accepted 31 January 2018

### KEYWORDS

aquaporins; osmo-sensing; volume regulation; volume-sensitive ion channels; transient receptor potential vanilloid 4 channel; TRPV4

### Introduction

Maintenance of cell volume is a homeostatic imperative for cells of most origins [1] and cellular responses suited to restore cell volume are set in motion in response to conditions causing cell swelling or shrinkage. Despite the widespread phenomenon and the numerous situations in which cell and tissue volume is compromised during physiology and pathophysiology, the molecular mechanisms governing cell volume homeostasis remain disputed and/or undefined. While many of the suggested volume regulatory (transport) mechanisms appear to sense volume changes, rather than aberrant changes in surrounding osmotic pressure, a few bona-fide osmo-sensing channels and transporters, i.e. BetP, ProP and OpuA have been discovered [2–6]. These membrane proteins sense changes in the osmolarity of the surrounding fluid via alterations in membrane properties/protein-lipid interactions and/or by changes in the intracellular  $K^+$  concentration, although direct effects via changes in the hydration state of the proteins cannot be excluded [7]. Despite the fact that abruptly arising, large osmotic challenges rarely, if ever, occur within mammalian tissue,

a convenient, and therefore common, manner of investigating volume regulatory responses is by introduction of excessive non-physiological osmotic challenges of up to 250 mOsm [8–16]. It is, in addition, not unusual that hyposmotic challenges of this magnitude are experimentally obtained by a dilution of the physiological test solution with distilled water [13–15,17]. Although this experimental approach is intended to simply change the osmotic pressure, the ionic strength will be diminished. This dilution thus causes drastic changes within the ionic equilibrium potentials, associated with a shift in the membrane potential and, in addition, the driving forces for a range of ion channels and transporters. These side effects are predicted to introduce serious confounding elements into the experimental set-up and may thus skew the outcome. We recommend that mannitol is included in the control solution (equiosmolar replacement of NaCl) and its subsequent removal will thus reduce the osmolarity without affecting the ionic strength [9,18,19].

A range of studies imply a requirement for aquaporins (AQPs) for activation of volume-sensitive ion channels,

**CONTACT** Associate Professor Nanna MacAulay  [macaulay@sund.ku.dk](mailto:macaulay@sund.ku.dk)  Department of Neuroscience, University of Copenhagen, Nørre Allé 14, 2200 Copenhagen N, Denmark.


Addendum to: Trine L. Toft-Bertelsen, David Krizaj and Nanna MacAulay. When size matters: Transient receptor potential vanilloid 4 channel as a volume sensor. *J Physiol* 2017; 595(11):3287–3302.

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.  
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

and possibly regulatory events, to take place [10,20,21] (Fig. 1). However, as AQPs are required to translate an abruptly introduced osmotic challenge into swift cell volume changes, the non-physiological experimental approach of introducing large osmotic gradients favors such conclusions [8,9,19,22]. In most physiological settings, the buildup of osmotic gradients will necessarily occur gradually with increased transmembrane transport activity and will represent the rate-limiting factor for cell swelling – rather than the osmotic water permeability (and hence AQP expression) of the cell membrane. Of interest, with a few exceptions such as pyramidal neurons [23] and stomach epithelium, the majority of cellular structures are rather water permeable even in the absence of AQPs, most likely via the multitude of water permeable cotransporters and uniporters present in these cells [24–26]. Striking examples are the AQP4-expressing astrocytes, the water permeability of which is reduced by only 50% (at 37°C) upon AQP4 knock-out [15] or knock-down [16]. As these cells are extremely water permeable to begin with, a 50% reduction of their osmotic water permeability still leaves them with the ability to equilibrate with a sudden osmotic challenge of  $-140$  mOsm well within a few seconds, at the physiological temperature of 37°C (Fig. 2). It follows that a physiologically relevant gradual buildup of an osmotic gradient (e.g. 1 mOsm at a time) via altered transport/channel activity will be instantly equilibrated – whether AQP4 is expressed or not. This phenomenon was illustrated in an experimental setting with astrocytic cell swelling in the absence of an introduced osmotic challenge: During neuronal activity in murine hippocampal slices (achieved by electric stimulation), the extracellular space (ECS) shrinks, mainly due to astrocytic cell swelling [27,28]. Genetic deletion of AQP4 did not prevent, or even reduce, this cell swelling [21], illustrating that the osmotic water permeability of the astrocytic cell membrane does not represent the rate-limiting step during astrocytic cell swelling (at least under these experimental conditions). We later demonstrated that the stimulus-induced astrocytic cell swelling relies on cotransporter-mediated water transport [27].


### Cell volume regulation

For cells to maintain their volume when faced with a stressor causing alterations in cell volume, their cellular response depends on a volume sensor coupled to a downstream mechanism causing transmembrane shift of



**Figure 1.** Contribution of AQP4 to astrocytic swelling and  $\text{Ca}^{2+}$  signaling. A: Cortical slices from wildtype (filled symbols) and AQP4-deficient (open symbols) mice were exposed to a hyposmotic gradient ( $-20\%$ , 60 mOsm (dashed lines)). Such osmotic challenge induced astrocytic swelling in slices from wildtype mice while these were non-detectable in slices from AQP4-deficient mice (marked with a red box), monitored with two-photon imaging of slices loaded with Texas red hydrazide. Verification of selective dye uptake in astrocytes was confirmed by two-photon imaging of slices from mice expressing GFP in astrocytes (*Glt-1-EGFP* BAC transgenic mice). A larger gradient of  $-30\%$ , 90 mOsm (solid lines) induced similar swelling in slices from both strains of mice. B: Astrocytes from wildtype (top panel) and AQP4-deficient (lower panel) mice were loaded with Rhod2 AM and exposed to a hyposmotic gradient of  $-20\%$  mOsm.  $\text{Ca}^{2+}$  dynamics were monitored  $\sim 200$  sec after introduction of the osmotic challenge (and the equivalent swelling marked with the red box in panel A). At this time point, wildtype astrocytes displayed  $\text{Ca}^{2+}$  spikes while AQP4-deficient astrocytes did not, presumably due to their absent (or limited) cell swelling. Modified from [20] with permission.

osmotic particles (either as separate entities or residing within one molecular transducer). Volume regulation most likely does not rely on one set of mechanisms but is rather predicted to depend on the molecular make-up of



**Figure 2.** Characterization of water transport in astrocytes from wild-type and AQP4-deficient mice. Primary culture of astrocytes from wildtype mice (top panel) or AQP4-deficient mice (lower panel) were faced with a hyposmotic challenge of  $-140$  mOsm (marked with an arrow) as obtained by 1:1 dilution of PBS with distilled water. The cellular swelling was monitored at 12°C (left) or 37°C (right) by the calcein quenching method. Modified from [15] with permission.

each cell type. A range of cotransport mechanisms have been indicated in cell volume regulation, with the  $\text{Na}^+/\text{K}^+/\text{2Cl}^-$  cotransporter and the  $\text{Na}^+/\text{H}^+$  exchanger as prominent inducers of cell swelling following osmotically induced cell shrinkage and, oppositely, the  $\text{K}^+/\text{Cl}^-$  cotransporter as an inducer of cell shrinkage upon osmotically induced cell swelling (for review, see [29–32]). An intricate regulatory pathway involving cell volume-sensitive kinases of the SPAK/WNK families leads to (de)phosphorylation of the abovementioned transporters, which in turn activates or inhibits the relevant set of transporters [33,34] (for review, see [31]) to promote the appropriate regulatory volume response. These kinases may be sensitive to cell volume changes themselves and/or to the altered intracellular  $\text{Cl}^-$  concentration coming about by the transmembrane shift of water during a large osmotic challenge [35]. An alternative mechanism involved in acute volume regulation following osmotically-induced cell swelling involves a concerted activity of plasma membrane  $\text{K}^+$  and  $\text{Cl}^-$  channels. This channel activity results in loss of KCl across the plasma membrane with a proposed associated directional water flow. It must be emphasized that this paradigm exclusively will work provided that both conductive pathways coexist and the electrochemical driving forces promote efflux of both ions: The principle of electroneutrality states that an equal amount of positive and negative charges must exist at either side of the membrane (with a miniscule excess of negative charges intracellularly to provide the negative membrane potential), as the lipid bilayer cannot sustain the large voltages that would occur given a millimolar excess of either positive or negative charges on one side of the membrane [36]. It is thus inconceivable that a buildup of an osmotic gradient of several mOsm (required for osmotically-induced water transport) could occur by the activity of a  $\text{K}^+$  (or  $\text{Cl}^-$  or  $\text{Na}^+$ ) channel alone, although promoted as a feasible manner of driving water movement [37–41]. In the central nervous system, release of several millimolar  $\text{K}^+$  into the ECS (to provide the required driving force for osmotic water efflux) would lead to neuronal depolarization and enhanced excitability. Such a mechanism may thus not be a viable manner of cell volume regulation in the cell structures of the central nervous system.

The activity of a range of different ion channels is modulated by altered cell volume, prominently so both  $\text{K}^+$  channels [42–45] and  $\text{Cl}^-$  channels [46–50] (extensively reviewed in [51,52]). This swelling-induced activation of  $\text{K}^+$  and  $\text{Cl}^-$  channels can occur as a direct sensing

of altered cell volume or, alternatively, secondarily to swelling-induced intracellular  $\text{Ca}^{2+}$  transients. Such  $\text{Ca}^{2+}$  dynamics may lead to activation of ion channels such as  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  channels [53,54] and/or initiation of cell signaling pathways with downstream activation of other transport mechanisms effectuating cell volume regulation. However, the linkage between cellular modulators that cause the volume-mediated  $\text{Ca}^{2+}$  dynamics and molecular pathways effectuating the volume regulatory response remains unresolved. Several volume-/stretch-sensitive  $\text{Ca}^{2+}$ -permeable channels have been identified [55–61], one of which is the transient receptor potential vanilloid 4 (TRPV4) ion channel. TRPV4, also known as the vanilloid receptor related and osmotically activated channel (VR-OAC [62]) or the OSM9-like transient receptor potential channel, member 4 (OTRPC4 [62,63]), is a nonselective cation channel expressed in many cell types: i.e. Müller cells, astrocytes, in neurons of the circumventricular organs, in mechanosensitive neurons of the mammalian inner ear hair cells, in the epithelia of the trachea, oviduct, lung and cochlea, in the airway, blood vessels, and in certain segments of the kidneys [62–70], many of which are key mechanosensory and osmosensory tissues.

### TRPV4 – an ion channel sensitive to abrupt volume changes

TRPV4 is a cation-permeable ion channel originally described as an osmo-sensor due to its activation by large hyposmotic challenges [63,71]. With such externally applied osmotic challenges, co-expression of an(y) AQP promotes the fast rate of osmotically-induced cell swelling leading to TRPV4 activation in astrocytes, Müller cells, and TRPV4-expressing oocytes [8,9,19,22]. However, TRPV4 activation readily occurred by cell swelling induced by cotransport of water *in the absence* of an externally applied osmotic challenge and presence of an AQP [9] and TRPV4 is therefore a genuine sensor of abrupt volume changes irrespective of the origin of the cell swelling. This observation thus promotes the term ‘volume-sensitive’ rather than ‘osmo-sensitive’ [9]. Nevertheless, the structural determinant underlying the repeatedly observed coupling of cell swelling to TRPV4 channel opening [8–11,19,22,62,63] remains unresolved.


With its robust activation upon cell volume increase, TRPV4 has been proposed as a candidate to translate cell swelling to cell volume regulation



[10,19,54,72–78] via its  $\text{Ca}^{2+}$  permeability, which places it in a favorable position as a main conductor of the  $\text{Ca}^{2+}$  influx required for volume regulatory mechanisms in some settings. The studies addressing the role of TRPV4 in regulatory volume responses were all performed by introduction of a non-physiologically large hyposmotic challenge, which is likely to introduce experimental confounders that may complicate the identification of the volume sensor and/or the molecular mechanisms effectuating the regulatory volume decrease. Of note, as TRPV4-mediated  $\text{Ca}^{2+}$  influx, in itself, cannot generate regulatory volume decrease, TRPV4 activity must necessarily couple to downstream effectors, currently unidentified, which then promote the loss of electrolytes and water.

### Contribution of TRPV4 to cell volume regulation during physiological relevant events

To investigate a potential role for TRPV4 in regulating cell volume *during a physiologically relevant cell swelling in situ*, independent on an applied osmotic gradient, we approximated a native setting by using acute hippocampal brain slices from rats. Neuronal activity in the brain causes release of  $\text{K}^+$  into the ECS where astrocytes act as  $\text{K}^+$  sinks to prevent extracellular  $\text{K}^+$  accumulation and a subsequent widespread depolarization [28,79–82]. In parallel with the management of  $\text{K}^+$ , the ECS shrinks [27,28,83–85], a phenomenon primarily attributed to the swelling of nearby astrocytic structures [83,85–86]. Thus electrical stimulation resulting in synaptic activity leads to a brief change in cell volume *without application of an osmotic gradient to the test solution*. This experimental scenario allowed us to test the involvement of TRPV4 in the return of astrocytic cell volume following the stimulus-evoked cell swelling. The relative change in size of the ECS was monitored via a tetramethyl ammonium ( $\text{TMA}^+$ )-sensitive microelectrode upon bath application of  $\text{TMA}^+$  (1.5 mM) as described in detail in [27,28]. In brief: Electrical stimulation of the CA1 Schaeffer collaterals of the hippocampus results in a shrinkage of the ECS and thus an increase in concentration of the membrane impermeable  $\text{TMA}^+$ . This volume trace then serves as an indirect readout of (astrocytic) cell swelling; see representative traces of stimulus-evoked ECS shrinkage illustrated in (Fig. 3A–C, left). To resolve the involvement of TRPV4 in the volume response, the non-selective TRPV4 inhibitor ruthenium red (RR; 1  $\mu\text{M}$ ) was applied to the recording chamber and the volume response recorded after



**Figure 3.** TRPV4 plays a minor, if any, role in regulating astrocyte volume during stimulus-evoked ECS shrinkage in acute hippocampal slices from rats. Ion-sensitive microelectrodes were employed to measure the relative size of the ECS upon addition of 1.5 mM  $\text{TMA}^+$  to the test solution. A–C, left: Representative recordings of the relative ECS obtained in CA1 (stratum radiatum) upon electrical stimulation (20 Hz; indicated by black bar) from slices treated with RR (A), HC067047 (B) or GSK101 (C). The black traces illustrate the control condition and the colored traces the one obtained in the presence of either antagonists or agonist. A–C, right: Normalized data on the peak amplitude, and decay constants. D: Representative Western blot illustrating the expression of TRPV4 in whole hippocampal tissue homogenate (20  $\mu\text{g}$  total protein/lane), in an astrocyte-enriched fraction of hippocampus (20  $\mu\text{g}$  total protein/lane), and in TRPV4-expressing- and uninjected *Xenopus* oocytes (amount of protein corresponding to 0.5 oocyte). The analysis was performed with two sets of individual samples all run as duplicates. Data presented as mean  $\pm$  SEM and statistical significance determined with a Student's paired t-test. \*,  $p < 0.05$ ; ns; not significant.

~6 min (Fig. 3A). An additional volume response was recorded after another 10 min of inhibitor incubation to ensure that the inhibitor had reached its full effect at the time of recording (for all test conditions). Inhibition of TRPV4 caused a slight increase in peak ECS shrinkage (to  $105.5 \pm 1.4\%$  of control,  $p < 0.05$ ,  $n = 7$  slices from 4 rats) (Fig. 3A, right) but did not affect the regulatory

volume decrease as the rate of return to baseline volume was undisturbed (the decay constant was  $104.7 \pm 2.4\%$  of control,  $p = 0.10$ ), see (Fig. 3A, right). In a similar fashion, the more specific TRPV4 inhibitor HC067047 ( $1 \mu\text{M}$ ) affected the peak ECS shrinkage ( $116.9 \pm 4.7\%$  of control,  $p < 0.05$ ,  $n = 5$  slices from 3 rats), (Fig. 3B, right) but did not influence the return of the ECS back to baseline as evaluated from the decay constant ( $106.5 \pm 7.5\%$  of control,  $p = 0.43$ ) (Fig. 3B, right). Activation of TRPV4 with the agonist GSK1016790A (GSK101;  $100 \text{ nM}$ , Fig. 3C) had no significant effect on the ECS dynamics in hippocampal brain slices (peak amplitude amounting to  $115.0 \pm 6.4\%$  of control,  $p = 0.13$  (Fig. 3C, right), and the decay constant to  $107.7 \pm 9.5\%$ ,  $n = 6$  slices from 4 rats (Fig. 3C, right)). Taken together, there appears to be little, if any, effect on astrocytic regulatory volume mechanisms of either blocking or activating TRPV4 during stimulus-evoked ECS shrinkage in acute hippocampal slices from rats. Notably, during the experimental paradigm, no osmotic gradients were introduced and the induced cell volume change occurred solely as the result of synaptic activity. TRPV4 expression in hippocampal astrocytes was previously demonstrated by immunocytochemistry [11,87–92]. However, the cortical TRPV4 expression was later shown confined to a subset ( $\sim 30\%$ ) of astrocytes [93] and the astrocytic TRPV4 transcript level reported to be fairly low [94,95]. To determine TRPV4 expression in rat hippocampus, we performed Western blotting analysis on homogenates obtained from rat hippocampi (P21) and from astrocyte-enriched fractions (P20–23) [96]. Efficient antibody recognition was verified in membrane preparations from uninjected and rat TRPV4-expressing *Xenopus laevis* oocytes [9], and GAPDH was used as loading control. Although we did not detect TRPV4 expression in  $20 \mu\text{g}$  loaded protein (Fig. 3G), another research group detected hippocampal TRPV4 expression by Western blot upon loading a 4-fold larger quantity of lysate [88]. The discrepancy between the data obtained by transcriptomics, Western blot, and immunohistochemistry may indicate a low expression of TRPV4 in astrocytes *in situ* and/or in a select subset of astrocytes, as detected in cortex [93].

## Conclusion

Volume regulation is an essential property of all cell types and a range of volume-sensing ion channels and transporters have been revealed and implicated in cell volume regulation throughout the years. Nevertheless,

the finer details regarding *exactly* how volume changes occur in physiology, and which mechanisms are involved in returning cell volume to its origin, remain largely unresolved. TRPV4 is a well-established sensor of volume changes, although predominantly during swift cell swelling [9]. Regardless, its implication in the subsequent cell volume regulatory events is less evident, and possibly not even pertinent under more physiological experimental conditions, in which cell swelling is obtained in the absence of externally applied osmotic gradients. For reasons of technical ease, experimenters often introduce such non-physiologically large osmotic gradients to study the ensuing volume-dependent activity of transporters and ion channels. While cell swelling/shrinkage is readily obtained in this manner, mammalian cells hardly experience these conditions *in situ* and we may simply be studying a different rate-limiting step under such circumstances. Cells exposed to an osmotic gradient ought to swell to a similar degree, whether or not an AQP is expressed, and whether or not the osmotic gradient is applied externally or from altered cellular transport activity *in vivo*. However, *the rate* with which it swells will depend on expression of an AQP. Abrupt introduction of a large osmotic gradient will thus allow investigation of cellular responses to a swift cell swelling of a magnitude and speed rarely observed in physiology and most prominently in settings with expression of an AQP. As a range of pathologies are associated with altered fluid dynamics, it is imperative that the research field as a whole obtains better tools as well as understanding of the molecular mechanisms underlying these fascinating volume regulatory properties. Our aim with this follow-up article is to pinpoint a few potential pit-falls along the path to reach this understanding.

## Abbreviations

AQP4	aquaporin 4
ECS	extracellular space
GAPDH	Glyceraldehyde 3-phosphate dehydrogenase
GSK101	GSK1016790A
OTRPC4	OSM9-like transient receptor potential channel, member 4
RR	ruthenium red
SPAK/WNK	SPS1-related proline/alanine-rich kinase/with no lysine
TMA <sup>+</sup>	tetramethyl ammonium

**TRPV4** Transient receptor potential vanilloid 4  
**VR-OAC** vanilloid receptor related and osmotically activated channel

## Acknowledgments

We are grateful for the technical assistance provided by Anca Stoica.

## Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed


## Funding

This project was supported by the VELUX Foundation [VELUX33842] (TLTB), the Lundbeck Foundation [R151-2013-14120] (BRL), and the Novo Nordisk Foundation [NNF15OC0017052] (BRL).

## ORCID

Trine L. Toft-Bertelsen  <http://orcid.org/0000-0001-9009-0094>

Brian R. Larsen  <http://orcid.org/0000-0003-2654-6645>

Nanna MacAulay  <http://orcid.org/0000-0002-7800-6600>

## References

- [1] Okada Y, Maeno E. Apoptosis, cell volume regulation and volume-regulatory chloride channels. Comparative Biochem Physiol part A, Mol Integrative Physiol. **2001**;130(3):377–83. doi:10.1016/S1095-6433(01)00424-X.
- [2] Rubenhagen R, Ronsch H, Jung H, et al. Osmosensor and osmoregulator properties of the betaine carrier BetP from *Corynebacterium glutamicum* in proteoliposomes. J Biol Chem. **2000**;275(2):735–41. doi:10.1074/jbc.275.2.735.
- [3] van der Heide T, Stuart MC, Poolman B. On the osmotic signal and osmosensing mechanism of an ABC transport system for glycine betaine. EMBO J. **2001**;20(24):7022–32. doi:10.1093/emboj/20.24.7022.
- [4] Schiller D, Ott V, Kramer R, et al. Influence of membrane composition on osmosensing by the betaine carrier BetP from *Corynebacterium glutamicum*. J Biol Chem. **2006**;281(12):7737–46. doi:10.1074/jbc.M513052200.
- [5] Rubenhagen R, Morbach S, Kramer R. The osmoreactive betaine carrier BetP from *Corynebacterium glutamicum* is a sensor for cytoplasmic K<sup>+</sup>. EMBO J. **2001**;20(19):5412–20. doi:10.1093/emboj/20.19.5412.
- [6] Schiller D, Kramer R, Morbach S. Cation specificity of osmosensing by the betaine carrier BetP of *Corynebacterium glutamicum*. FEBS Letters. **2004**;563(1-3):108–12. doi:10.1016/S0014-5793(04)00279-0.
- [7] Wood JM. Osmosensing by bacteria. Sci STKE. **2006**;2006(357):pe43. doi:10.1126/stke.3572006pe43.
- [8] Ryskamp DA, Jo AO, Frye AM, et al. Swelling and eicosanoid metabolites differentially gate TRPV4 channels in retinal neurons and glia. J Neurosci Official J Soc Neurosci. **2014**;34(47):15689–700. doi:10.1523/JNEUROSCI.2540-14.2014.
- [9] Toft-Bertelsen TL, Krizaj D, MacAulay N. When size matters: transient receptor potential vanilloid 4 channel as a volume-sensor rather than an osmo-sensor. J Physiol. **2017**;595(11):3287–302. doi:10.1113/JP274135.
- [10] Benfenati V, Caprini M, Dovizio M, et al. An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. Proc Natl Acad Sci U S A. **2011**;108(6):2563–8. doi:10.1073/pnas.1012867108.
- [11] Benfenati V, Amiry-Moghaddam M, Caprini M, et al. Expression and functional characterization of transient receptor potential vanilloid-related channel 4 (TRPV4) in rat cortical astrocytes. Neuroscience. **2007**;148(4):876–92. doi:10.1016/j.neuroscience.2007.06.039.
- [12] Loukin S, Su Z, Kung C. Increased basal activity is a key determinant in the severity of human skeletal dysplasia caused by TRPV4 mutations. PloS One. **2011**;6(5):e19533. doi:10.1371/journal.pone.0019533.
- [13] Pannicke T, Iandiev I, Wurm A, et al. Diabetes alters osmotic swelling characteristics and membrane conductance of glial cells in rat retina. Diabetes. **2006**;55(3):633–9. doi:10.2337/diabetes.55.03.06.db05-1349.
- [14] Pannicke T, Iandiev I, Uckermann O, et al. A potassium channel-linked mechanism of glial cell swelling in the postischemic retina. Mol Cell Neurosci. **2004**;26(4):493–502. doi:10.1016/j.mcn.2004.04.005.
- [15] Solenov E, Watanabe H, Manley GT, et al. Sevenfold-reduced osmotic water permeability in primary astrocyte cultures from AQP-4-deficient mice, measured by a fluorescence quenching method. Am J Physiol Cell Physiol. **2004**;286(2):C426–32. doi:10.1152/ajpcell.00298.2003.
- [16] Nicchia GP, Frigeri A, Liuzzi GM, et al. Inhibition of aquaporin-4 expression in astrocytes by RNAi determines alteration in cell morphology, growth, and water transport and induces changes in ischemia-related genes. FASEB J Official Publication Federation Am Soc Exp Biol. **2003**;17(11):1508–10. doi:10.1096/fj.02-1183fj.
- [17] Vardjan N, Horvat A, Anderson JE, et al. Adrenergic activation attenuates astrocyte swelling induced by hypotonicity and neurotrauma. Glia. **2016**;64(6):1034–49.
- [18] Steffensen AB, Sword J, Croom D, et al. Chloride Cotransporters as a Molecular Mechanism underlying Spreading Depolarization-Induced Dendritic Beading. The J Neurosci Official J Soc Neurosci. **2015**;35(35):12172–87. doi:10.1523/JNEUROSCI.0400-15.2015.
- [19] Jo AO, Ryskamp DA, Phuong TT, et al. TRPV4 and AQP4 channels synergistically regulate cell volume and calcium homeostasis in retinal Muller Glia. J neurosci Official J Soc Neurosci. **2015**;35(39):13525–37. doi:10.1523/JNEUROSCI.1987-15.2015.
- [20] Thrane AS, Rappold PM, Fujita T, et al. Critical role of aquaporin-4 (AQP4) in astrocytic Ca<sup>2+</sup> signaling events elicited by cerebral edema. Proc Natl Acad Sci U S A. **2011**;108(2):846–51. doi:10.1073/pnas.1015217108.



- [21] Haj-Yasein NN, Jensen V, Ostby I, et al. Aquaporin-4 regulates extracellular space volumedynamics during high-frequency synaptic stimulation: a gene deletion study in mouse hippocampus. *Glia*. 2012;60(6):867–74. doi:10.1002/glia.22319.
- [22] Mola MG, Sparaneo A, Gargano CD, et al. The speed of swelling kinetics modulates cell volume regulation and calcium signaling in astrocytes: A different point of view on the role of aquaporins. *Glia*. 2016;64(1):139–54. doi:10.1002/glia.22921.
- [23] Andrew RD, Labron MW, Boehnke SE, et al. Physiological evidence that pyramidal neurons lack functional water channels. *Cereb Cortex*. 2007;17(4):787–802. doi:10.1093/cercor/bhk032.
- [24] MacAulay N, Hamann S, Zeuthen T. Water transport in the brain: role of cotransporters. *Neuroscience*. 2004;129(4):1031–44. doi:10.1016/j.neuroscience.2004.06.045.
- [25] MacAulay N, Zeuthen T. Water transport between CNS compartments: contributions of aquaporins and cotransporters. *Neuroscience*. 2010;168(4):941–56. doi:10.1016/j.neuroscience.2009.09.016.
- [26] Macaulay N, Zeuthen T. Glial K(+) clearance and cell swelling: key roles for cotransporters and pumps. *Neurochem Res*. 2012;37(11):2299–309. doi:10.1007/s11064-012-0731-3.
- [27] Larsen BR, MacAulay N. Activity-dependent astrocyte swelling is mediated by pH-regulating mechanisms. *Glia*. 2017;65(10):1668–81. doi:10.1002/glia.23187.
- [28] Larsen BR, Assentoft M, Cotrina ML, et al. Contributions of the Na(+)/K(+)-ATPase, NKCC1, and Kir4.1 to hippocampal K(+) clearance and volume responses. *Glia*. 2014;62(4):608–22. doi:10.1002/glia.22629.
- [29] Mongin AA, Orlov SN. Mechanisms of cell volume regulation and possible nature of the cell volume sensor. *Pathophysiology*. 2001;8(2):77–88. doi:10.1016/S0928-4680(01)00074-8.
- [30] Gamba G. Molecular physiology and pathophysiology of electroneutral cation-chloride cotransporters. *Physiological Rev*. 2005;85(2):423–93. doi:10.1152/physrev.00011.2004.
- [31] Kahle KT, Khanna AR, Alper SL, et al. K-Cl cotransporters, cell volume homeostasis, and neurological disease. *Trends Mol Med*. 2015;21(8):513–23. doi:10.1016/j.molmed.2015.05.008.
- [32] Pedersen SF, O'Donnell ME, Anderson SE, et al. Physiology and pathophysiology of Na<sup>+</sup>/H<sup>+</sup> exchange and Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> cotransport in the heart, brain, and blood. *Am J Physiol Regulatory Integrative Comparative Physiol*. 2006;291(1):R1–25. doi:10.1152/ajpregu.00782.2005.
- [33] Kahle KT, Rinehart J, Lifton RP. Phosphoregulation of the Na-K-2Cl and K-Cl cotransporters by the WNK kinases. *Biochim Et Biophys Acta*. 2010;1802(12):1150–8. doi:10.1016/j.bbadis.2010.07.009.
- [34] de Los Heros P, Kahle KT, Rinehart J, et al. WNK3 bypasses the tonicity requirement for K-Cl cotransporter activation via a phosphatase-dependent pathway. *Proc Natl Acad Sci U S A*. 2006;103(6):1976–81. doi:10.1073/pnas.0510947103.
- [35] Zhang J, Gao G, Begum G, et al. Functional kinomics establishes a critical node of volume-sensitive cation-Cl<sup>-</sup> cotransporter regulation in the mammalian brain. *Scientific Reports*. 2016;6:35986. doi:10.1038/srep35986.
- [36] Blaustein MP, Kao JP, Matteson DR. Cellular physiology and neurophysiology E-Book: Mosby physiology monograph series, The Curtis Center, Philadelphia, PA: Elsevier Health Sciences; 2011.
- [37] Stokum JA, Kwon MS, Woo SK, et al. SUR1-TRPM4 and AQP4 form a heteromultimeric complex that amplifies ion/water osmotic coupling and drives astrocyte swelling. *Glia*. 2018;66(1):108–25. doi:10.1002/glia.23231.
- [38] Hoffmann EK, Lambert IH, Pedersen SF. Physiology of cell volume regulation in vertebrates. *Physiol Rev*. 2009;89(1):193–277. doi:10.1152/physrev.00037.2007.
- [39] Nagelhus EA, Mathiesen TM, Ottersen OP. Aquaporin-4 in the central nervous system: cellular and subcellular distribution and coexpression with KIR4.1. *Neuroscience*. 2004;129(4):905–13. doi:10.1016/j.neuroscience.2004.08.053.
- [40] Kimelberg HK, Macvicar BA, Sontheimer H. Anion channels in astrocytes: biophysics, pharmacology, and function. *Glia*. 2006;54(7):747–57. doi:10.1002/glia.20423.
- [41] Rungta RL, Choi HB, Tyson JR, et al. The cellular mechanisms of neuronal swelling underlying cytotoxic edema. *Cell*. 2015;161(3):610–21. doi:10.1016/j.cell.2015.03.029.
- [42] Tejada MA, Hashem N, Calloe K, et al. Heteromeric Slick/Slack K<sup>+</sup> channels show graded sensitivity to cell volume changes. *PloS One*. 2017;12(2):e0169914. doi:10.1371/journal.pone.0169914.
- [43] Calloe K, Elmedyb P, Olesen SP, et al. Hypoosmotic cell swelling as a novel mechanism for modulation of cloned HCN2 channels. *Biophysical J*. 2005;89(3):2159–69. doi:10.1529/biophysj.105.063792.
- [44] Grunnet M, MacAulay N, Jorgensen NK, et al. Regulation of cloned, Ca<sup>2+</sup>-activated K<sup>+</sup> channels by cell volume changes. *Pflugers Archiv: Eur J Physiol*. 2002;444(1-2):167–77. doi:10.1007/s00424-002-0782-4.
- [45] Grunnet M, Jespersen T, MacAulay N, et al. KCNQ1 channels sense small changes in cell volume. *J Physiol*. 2003;549(Pt 2):419–27. doi:10.1113/jphysiol.2003.038455.
- [46] Pedersen SF, Okada Y, Nilius B. Biophysics and Physiology of the Volume-Regulated Anion Channel (VRAC)/Volume-Sensitive Outwardly Rectifying Anion Channel (VSOR). *Pflugers Archiv: Eur J Physiol*. 2016;468(3):371–83. doi:10.1007/s00424-015-1781-6.
- [47] Nilius B, Eggermont J, Voets T, et al. Properties of volume-regulated anion channels in mammalian cells. *Progress Biophys Mol Biol*. 1997;68(1):69–119. doi:10.1016/S0079-6107(97)00021-7.
- [48] Cahalan MD, Lewis RS. Role of potassium and chloride channels in volume regulation by T lymphocytes. *Soc Gen Physiol Ser*. 1988;43:281–301.
- [49] Schober AL, Wilson CS, Mongin AA. Molecular composition and heterogeneity of the LRRC8-containing swelling-activated osmolyte channels in primary rat astrocytes. *J Physiol*. 2017;595(22):6939–51. doi:10.1113/JP275053.

- [50] Voss FK, Ullrich F, Munch J, et al. Identification of LRRC8 heteromers as an essential component of the volume-regulated anion channel VRAC. *Science*. **2014**;344(6184):634–8. doi:10.1126/science.1252826.
- [51] Mongin AA. Volume-regulated anion channel—a frenemy within the brain. *Pflugers Archiv: Eur J Physiol*. **2016**;468(3):421–41. doi:10.1007/s00424-015-1765-6.
- [52] Jentsch TJ. VRACs and other ion channels and transporters in the regulation of cell volume and beyond. *Nat Rev Mol Cell Biol*. **2016**;17(5):293–307. doi:10.1038/nrm.2016.29.
- [53] Nilius B, Watanabe H, Vriens J. The TRPV4 channel: structure-function relationship and promiscuous gating behaviour. *Pflugers Archiv: Eur J Physiol*. **2003**;446(3):298–303. doi:10.1007/s00424-003-1028-9.
- [54] Fernandez-Fernandez JM, Nobles M, Currid A, et al. Maxi K<sup>+</sup> channel mediates regulatory volume decrease response in a human bronchial epithelial cell line. *Am J Physiol Cell Physiol*. **2002**;283(6):C1705–14. doi:10.1152/ajpcell.00245.2002.
- [55] Zhang Z, Wen Y, Du J, et al. Effects of mechanical stretch on the functions of BK and L-type Ca(2<sup>+</sup>) channels in vascular smooth muscle cells. *J Biomech*. **2017**;67:18–23. doi:10.1016/j.jbiomech.2017.11.015.
- [56] McGahon MK, Fernandez JA, Dash DP, et al. TRPV2 Channels Contribute to Stretch-Activated Cation Currents and Myogenic Constriction in Retinal Arterioles. *Invest Ophthalmol Vis Sci*. **2016**;57(13):5637–47. doi:10.1167/iovs.16-20279.
- [57] Ciura S, Liedtke W, Bourque CW. Hypertonicity sensing in organum vasculosum lamina terminalis neurons: a mechanical process involving TRPV1 but not TRPV4. *J Neurosci: Official J Soc Neurosci*. **2011**;31(41):14669–76. doi:10.1523/JNEUROSCI.1420-11.2011.
- [58] Khajavi N, Reinach PS, Skrzypski M, et al. L-carnitine reduces in human conjunctival epithelial cells hypertonic-induced shrinkage through interacting with TRPV1 channels. *Cell Physiol Biochem Int J Exp Cell Physiol Biochem Pharmacol*. **2014**;34(3):790–803. doi:10.1159/000363043.
- [59] Maroto R, Raso A, Wood TG, et al. TRPC1 forms the stretch-activated cation channel in vertebrate cells. *Nat Cell Biol*. **2005**;7(2):179–85. doi:10.1038/ncb1218.
- [60] Aguetaz E, Bois P, Cognard C, et al. Stretch-activated TRPV2 channels: Role in mediating cardiopathies. *Progress Biophys Mol Biol*. **2017**;130(Pt B):273–80. doi:10.1016/j.pbiomolbio.2017.05.007.
- [61] Filipovic D, Sackin H. A calcium-permeable stretch-activated cation channel in renal proximal tubule. *Am J Physiol*. **1991**;260(1 Pt 2):F119–29.
- [62] Liedtke W, Choe Y, Marti-Renom MA, et al. Vanilloid receptor-related osmotically activated channel (VR-OAC), a candidate vertebrate osmoreceptor. *Cell*. **2000**;103(3):525–35. doi:10.1016/S0092-8674(00)00143-4.
- [63] Strotmann R, Harteneck C, Nunnenmacher K, et al. OTRPC4, a nonselective cation channel that confers sensitivity to extracellular osmolarity. *Nat Cell Biol*. **2000**;2(10):695–702. doi:10.1038/35036318.
- [64] Wissenbach U, Boddling M, Freichel M, et al. Trp12, a novel Trp related protein from kidney. *FEBS Letters*. **2000**;485(2-3):127–34. doi:10.1016/S0014-5793(00)02212-2.
- [65] Liedtke W, Friedman JM. Abnormal osmotic regulation in trpv4<sup>-/-</sup> mice. *Proc Natl Acad Sci U S A*. **2003**;100(23):13698–703. doi:10.1073/pnas.1735416100.
- [66] Jia Y, Wang X, Varty L, et al. Functional TRPV4 channels are expressed in human airway smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol*. **2004**;287(2):L272–8. doi:10.1152/ajplung.00393.2003.
- [67] Tian W, Salanova M, Xu H, et al. Renal expression of osmotically responsive cation channel TRPV4 is restricted to water-impermeant nephron segments. *Am J Physiol Renal Physiol*. **2004**;287(1):F17–24. doi:10.1152/ajprenal.00397.2003.
- [68] Andrade YN, Fernandes J, Vazquez E, et al. TRPV4 channel is involved in the coupling of fluid viscosity changes to epithelial ciliary activity. *J Cell Biol*. **2005**;168(6):869–74. doi:10.1083/jcb.200409070.
- [69] Earley S, Pauyo T, Drapp R, et al. TRPV4-dependent dilation of peripheral resistance arteries influences arterial pressure. *Am J Physiol Heart Circulatory Physiol*. **2009**;297(3):H1096–102. doi:10.1152/ajpheart.00241.2009.
- [70] Ryskamp DA, Witkovsky P, Barabas P, et al. The polymodal ion channel transient receptor potential vanilloid 4 modulates calcium flux, spiking rate, and apoptosis of mouse retinal ganglion cells. *J Neurosci Official J Soc Neurosci*. **2011**;31(19):7089–101. doi:10.1523/JNEUROSCI.0359-11.2011.
- [71] Liedtke W, Tobin DM, Bargmann CI, et al. Mammalian TRPV4 (VR-OAC) directs behavioral responses to osmotic and mechanical stimuli in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*. **2003**;100(Suppl 2):14531–6. doi:10.1073/pnas.2235619100.
- [72] Galizia L, Pizzoni A, Fernandez J, et al. Functional interaction between AQP2 and TRPV4 in renal cells. *J Cell Biochem*. **2012**;113(2):580–9. doi:10.1002/jcb.23382.
- [73] Pan Z, Yang H, Mergler S, et al. Dependence of regulatory volume decrease on transient receptor potential vanilloid 4 (TRPV4) expression in human corneal epithelial cells. *Cell Calcium*. **2008**;44(4):374–85. doi:10.1016/j.ceca.2008.01.008.
- [74] Becker D, Bereiter-Hahn J, Jendrach M. Functional interaction of the cation channel transient receptor potential vanilloid 4 (TRPV4) and actin in volume regulation. *Eur J Cell Biol*. **2009**;88(3):141–52. doi:10.1016/j.ejcb.2008.10.002.
- [75] Arniges M, Vazquez E, Fernandez-Fernandez JM, et al. Swelling-activated Ca<sup>2+</sup> entry via TRPV4 channel is defective in cystic fibrosis airway epithelia. *J Biol Chem*. **2004**;279(52):54062–8. doi:10.1074/jbc.M409708200.
- [76] Liu X, Bandyopadhyay BC, Nakamoto T, et al. A role for AQP5 in activation of TRPV4 by hypotonicity: concerted involvement of AQP5 and TRPV4 in regulation of cell volume recovery. *J Biol Chem*. **2006**;281(22):15485–95. doi:10.1074/jbc.M600549200.

- [77] Becker D, Blase C, Bereiter-Hahn J, et al. TRPV4 exhibits a functional role in cell-volume regulation. *J Cell Sci.* **2005**;118(Pt 11):2435–40. doi:10.1242/jcs.02372.
- [78] Jo AO, Lakk M, Frye AM, et al. Differential volume regulation and calcium signaling in two ciliary body cell types is subserved by TRPV4 channels. *Proc Natl Acad Sci U S A.* **2016**;113(14):3885–90. doi:10.1073/pnas.1515895113.
- [79] Hertz L, Song D, Xu J, et al. Role of the Astrocytic Na(+), K(+)-ATPase in K(+) Homeostasis in Brain: K(+) Uptake, Signaling Pathways and Substrate Utilization. *Neurochem Res.* **2015**;40(12):2505–16. doi:10.1007/s11064-014-1505-x.
- [80] Kofuji P, Newman EA. Potassium buffering in the central nervous system. *Neuroscience.* **2004**;129(4):1045–56. doi:10.1016/j.neuroscience.2004.06.008.
- [81] Walz W. Role of astrocytes in the clearance of excess extracellular potassium. *Neurochem Int.* **2000**;36(4-5):291–300. doi:10.1016/S0197-0186(99)00137-0.
- [82] Larsen BR, Stoica A, MacAulay N. Managing Brain Extracellular K(+) during Neuronal Activity: The Physiological Role of the Na(+)/K(+)-ATPase Subunit Isoforms. *Frontiers Physiol.* **2016**;7:141. doi:10.3389/fphys.2016.00141.
- [83] Connors BW, Ransom BR, Kunis DM, et al. Activity-dependent K+ accumulation in the developing rat optic nerve. *Science.* **1982**;216(4552):1341–3. doi:10.1126/science.7079771.
- [84] Dietzel I, Heinemann U, Hofmeier G, et al. Transient changes in the size of the extracellular space in the sensorimotor cortex of cats in relation to stimulus-induced changes in potassium concentration. *Exp Brain Res.* **1980**;40(4):432–9. doi:10.1007/BF00236151.
- [85] Ransom BR, Yamate CL, Connors BW. Activity-dependent shrinkage of extracellular space in rat optic nerve: a developmental study. *J Neurosci.* **1985**;5(2):532–5.
- [86] MacVicar BA, Feighan D, Brown A, et al. Intrinsic optical signals in the rat optic nerve: role for K(+) uptake via NKCC1 and swelling of astrocytes. *Glia.* **2002**;37(2):114–23. doi:10.1002/glia.10023.
- [87] Butenko O, Dzamba D, Benesova J, et al. The increased activity of TRPV4 channel in the astrocytes of the adult rat hippocampus after cerebral hypoxia/ischemia. *PloS One.* **2012**;7(6):e39959. doi:10.1371/journal.pone.0039959.
- [88] Bai JZ, Lipski J. Differential expression of TRPM2 and TRPV4 channels and their potential role in oxidative stress-induced cell death in organotypic hippocampal culture. *Neurotoxicology.* **2010**;31(2):204–14. doi:10.1016/j.neuro.2010.01.001.
- [89] Bai JZ, Lipski J. Involvement of TRPV4 channels in Abeta (40)-induced hippocampal cell death and astrocytic Ca(2+) signalling. *Neurotoxicology.* **2014**;41:64–72. doi:10.1016/j.neuro.2014.01.001.
- [90] Lipski J, Park TI, Li D, et al. Involvement of TRP-like channels in the acute ischemic response of hippocampal CA1 neurons in brain slices. *Brain Res.* **2006**;1077(1):187–99. doi:10.1016/j.brainres.2006.01.016.
- [91] Shi M, Du F, Liu Y, et al. Glial cell-expressed mechanosensitive channel TRPV4 mediates infrasound-induced neuronal impairment. *Acta Neuropathologica.* **2013**;126(5):725–39. doi:10.1007/s00401-013-1166-x.
- [92] Shibasaki K, Suzuki M, Mizuno A, et al. Effects of body temperature on neural activity in the hippocampus: regulation of resting membrane potentials by transient receptor potential vanilloid 4. *J Neurosci Official J Soc Neurosci.* **2007**;27(7):1566–75. doi:10.1523/JNEUROSCI.4284-06.2007.
- [93] Shibasaki K, Ikenaka K, Tamalu F, et al. A novel subtype of astrocytes expressing TRPV4 (transient receptor potential vanilloid 4) regulates neuronal excitability via release of gliotransmitters. *J Biol Chem.* **2014**;289(21):14470–80. doi:10.1074/jbc.M114.557132.
- [94] Chai H, Diaz-Castro B, Shigetomi E, et al. Neural Circuit-Specialized Astrocytes: Transcriptomic, Proteomic, Morphological, and Functional Evidence. *Neuron.* **2017**;95(3):531–49 e9. doi:10.1016/j.neuron.2017.06.029.
- [95] Zhang Y, Chen K, Sloan SA, et al. An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci Official J Soc Neurosci.* **2014**;34(36):11929–47. doi:10.1523/JNEUROSCI.1860-14.2014.
- [96] Stoica A, Larsen BR, Assentoft M, et al. The alpha2beta2 isoform combination dominates the astrocytic Na(+) /K(+) -ATPase activity and is rendered nonfunctional by the alpha2.G301R familial hemiplegic migraine type 2-associated mutation. *Glia.* **2017**;65(11):1777–93. doi:10.1002/glia.23194.